

TOTAL MENTAL SERVICE SERVICES

TOOLA SACCESSALTOO ON THE RESIDENT TOTAL TELEGRADISSINGS TOOL

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A



OFFICE OF NAVAL RESEARCH

Contract N00014-82-K-0113

Task No. NR 359-258

TECHNICAL REPORT NO. NYU/DC/TR-11-NEW SERIES-2


TRANSITION DIPOLE-SOLVENT INTERACTION IN PHOTOIONIZATION

by

Paul Delahay and Andrew Dziedzic

Accepted for publication in Chemical Physics Letters

New York University Department of Chemistry New York, NY



September 1986

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

86 10 17 004 86 10 17 004

	tressified	NACIONAL PARAMENTAL	THE PROPERTY OF THE PARTY OF			<u> Anthropologica (* 1</u>	ETATATATATATATATATATA	
	SIFICATION OF							
			REPORT DOCUM	MENTATION	PAGE			
a. REPORT SECURITY CLASSIFICATION				16. RESTRICTIVE MARKINGS				
	NA			NA NA				
a. SECURITY	CLASSIFICATION	N AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT This document has been approved for public				
NA  2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				Inis docume release and	nt mas be lsale; it	æen approved ts distribut	ion is unlimite	
	NA NA							
I. PERFORMING ORGANIZATION REPORT NUMBER(S)				5. MONITORING ORGANIZATION REPORT NUMBER(S)				
NYU/DC/TR-11-NEW SERIES-2				NA				
Sa. NAME OF PERFORMING ORGANIZATION 6b. OFFICE SYMBOL				7a. NAME OF MONITORING ORGANIZATION				
Department of Chemistry New York University			(If applicable) NA	ONR				
Sc. ADDRESS (	City, State, and	d ZIP Code)		7b. ADDRESS (City, State, and ZIP Code)				
New	York, NY	10003		800	800 N. Quincy Street			
				Arli	ington, V	A 22217		
	a. NAME OF FUNDING/SPONSORING 8b. OFFICE SYMBOL			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER				
ORGANIZA	ORGANIZATION (If applicable) NA				Contract N00014-82-K-0113			
Bc. ADDRESS (	City, State, and	J ZIP Code)		10. SOURCE OF FUNDING NUMBERS				
		·		PROGRAM	PROJECT	TASK	WORK UNIT	
				ELEMENT NO.	NO.	NO.	ACCESSION NO	
·	1 Constant				<u> </u>			
	ude Security C		: Interaction in	Photoionizat	tion			
114	112101011 0	10016-30116114	1110614001011	11100010111	,			
12. PERSONAL Pau	. AUTHOR(S)	and Andrew D		<del></del>	<del></del>	<del></del>		
13a, TYPE OF		13b. TIME CO		14 DATE OF REPO	ORT (Year, Mo	onth. Dav) 15. PA	AGE COUNT	
	hnical	FROM	το	14. DATE OF REPO 9/30/86			20 ***	
	NTARY NOTAT	-						
Acc	epted for	publication	in Chemical Phys	sics Letters				
17.	COSATI	CODES	18. SUBJECT TERMS (	Continue on reverse			block number)	
FIELD	GROUP SUB-GROUP Dielectric				Solut			
			Dispersion sp			ition dipole	•	
	1	1	I Dhotoemission		VIIV 🤄	pectroscopy		

19. ABSTRACT (Continue on reverse if necessary and identify by block number) The interaction between solvent molecules and transition dipoles results in a measurable nonequilibrium electronic contribution to the energetics of photoionization in the case of significant dielectric dispension of the solvent at the prevailing photon energy. Earlier theoretical work on the resulting dispersion shift is justified on the basis of transition dipole-solvent interaction. Additional experimental evidence is presented.

CONTRACTOR CONTRACTOR COSERS	21. ABSTRACT SECURITY CLASSIFICATION Unclassified
22a NAME OF RESPONSIBLE INDIVIDUAL Paul Delahay	22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL 212-598-2811

**DD FORM 1473, 84 MAR** 

PONNOS OF TOTAL STREET STREET STREET STREET STREET STREET STREET STREET STREET STREET

83 APR edition may be used until exhausted. All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

TRANSITION DIPOLE-SOLVENT INTERACTION IN PHOTOIONIZATION

Paul DELAHAY and Andrew DZIEDZIC

Department of Chemistry, New York University, New York, New York 10003, USA Received

The interaction between solvent molecules and transition dipoles results in a measurable <u>nonequilibrium</u> electronic contribution to the energetics of photoionization in the case of significant dielectric dispersion of the solvent at the prevailing photon energy. Earlier theoretical work on the resulting <u>dispersion shift</u> is justified on the basis of transition dipole—solvent interaction. Additional experimental evidence is presented.

### 1. Introduction

Spectroscopic transitions from one bound state to another bound state involve a transition dipole which oscillates at or very near the frequency of the incident radiation [1]. Such an oscillating transition dipole is also involved in transitions to the continuum [2], but there is, of course, in that case ultimate separation of the emitted electron from its parent atomic or molecular system. In solutions, the transition dipole interacts with surrounding solvent molecules in a process which is similar to solvation of a dipole or, in the limit of complete charge separation, to ionic solvation except that nuclear motion is not involved. The response of the solvent is determined by its <u>dielectric properties</u> at the frequency at which the transition dipole oscillates, that is, by the real and imaginary optical dielectric constant of the solvent at or near the frequency of radiation. The complex dielectric constant is a function of photon energy because of dispersion and consequently the energy for the transition dipole—solvent interaction varies with the energy of the incident radiation.



A-1

odes

In the absence of dispersion, the free energy for transition dipole-solvent interaction depends on the limiting value  $\varepsilon_{op}^0$  of the optical dielectric constant (in the near infrared range). This free energy of interaction for the dielectric constant  $\varepsilon_{op}^0$  is part of the difference in free energy between the initial and final states of the photoionization process. The <u>difference</u> between the free energies of transition dipole-solvent interaction for the dielectric constant at the prevailing photon energy and  $\varepsilon_{op}^0$  therefore represents a <u>nonequilibrium electronic</u> contribution to the free energy of photoionization. The expression <u>dispersion</u> shift is suggested for this nonequilibrium electronic contribution.

Two main questions arise about the dispersion shift: (i) What experimental evidence is there which supports the introduction of a dispersion shift depending on photon energy as part of the photoionization energy in liquids and solutions? (ii) How can the dispersion shift be calculated? Experimental evidence was presented in earlier paper [3] and theory was developed in [4]. Additional evidence is reported in the present Letter and the previous theory is given further conceptual support.

### 2. Experimental evidence for the dispersion shift

Strong support for the occurrence of transition dipole-solvent interaction is available from experiments on photoelectron emission by liquids and solutions [5]. In these experiments, normally incident VUV radiation strikes the liquid surface and emitted photoelectrons are collected on a planar grid electrode parallel to the liquid surface. An emission yield, Y, is calculated as the number of collected electrons per incident photon. In the absence of a dispersion effect, Y is proportional to  $(E-E_{\rm t})^2$ , where E is the photon energy and  $E_{\rm t}$  the threshold energy for the species being photoionized. This quadratic emission law holds at photon energies at least a few tenths of an

electronvolt higher than  $E_t$ . Therefore, a plot of  $Y^{1/2}$  against E should be linear, and the derivative  $dY^{1/2}/dE$  should be independent of photon energy in the range of validity of the quadratic law. These conclusions hold if dispersion of the solvent does not affect the variations of Y with E. This is the case, for instance, for photoelectron emission by liquid water ( $E_t$  = 10.06 eV) because the effect of dispersion on emission for this substance is nearly independent of photon energy in the range of quadratic extrapolation from 10 to 11 eV.

Emission spectra (Y against E) and plots of  $Y^{1/2}$  against E generally exhibit a fine structure, and the derivative  $dY^{1/2}/dE$  is not independent of E (fig. 1) above the threshold energy. This effect was first reported for various inorganic anions in aqueous solution [3] but it is of general nature. The functional dependence of  $dY^{1/2}/dE$  on E (dispersion spectrum) is essentially determined by the solvent and in a minor way by the nature of the substance being photoionized. This is conclusively shown by a series of experiments.

Firstly, dispersion spectra are not the result of an experimental artefact having its origin in the vacuum ultraviolet radiation source since practically the same dispersion spectrum is obtained from 7 to 9 eV (fig. 1) with two sources having significantly different spectral outputs (fig. 2). The difference between the spectra in fig. 1 above 9 eV is caused by the very low output of the argon source in that range and the large error in the normalization of the photon flux.

Secondly, dispersion spectra do not result from an experimental artefact having its origin in the monochromator and optical system since very different spectra are obtained for iodide solutions in water and glycerol (fig. 3). The dispersion spectrum of the iodide solution in glycerol above 9.0 eV is quite

similar to the spectrum for pure glycerol because of dominant emission by the solvent. In a separate experiment, water vapor was allowed to be absorbed by the iodide solution in glycerol in the emission chamber. The dispersion spectrum evolved toward the spectrum obtained with aqueous solutions of iodide, as the water was increasingly absorbed by the glycerol.

The dispersion spectrum for a given solvent is essentially independent of kinetic energy of quasifree electrons produced by photoionization. This was shown in ref. [3] for aqueous solutions of 17 inorganic anions with threshold energies ranging from 7.2 to 8.9 eV.

3. Theoretical considerations on the dispersion shift

Two <u>limiting</u> cases may be considered in the calculation of the dispersion shift:

(i) One assumes complete separation of the emitted electron from its parent ion or molecule in solution. This approach was developed recently [4], and a detailed calculation was performed of the dispersion shift. A discrete model of solvent molecules was used for the inner-sphere region around the photoionized species, whereas the outer-sphere region was treated in terms of a continuous medium. Both transparent and absorbing solvents were considered.

The dispersion shift was calculated on the <u>ad hoc</u> assumption that the ionic field changes on a time scale corresponding to the radiation frequency. The central idea of the present paper, namely that of transition dipole-solvent interaction, provides the <u>conceptual</u> justification for the calculation in [4]. Only the change in the ionic field from the initial to the final state of photoionization was considered in [4], and no attempt was made to follow the time-evolution of the transition dipole-solvent interaction. Thus, the matrix element for absorption refers only to the initial and final states, and the absorption of a photon is accounted for by the annihilation operator acting in occupation space [6].

The theory of [4] accounts very well for the functional dependence on photon energy exhibited by experimental dispersion spectra. In particular, excellent agreement was obtained between theory and experiment for the photon energies at the 12 extrema exhibited by dispersion spectra of aqueous solutions between 7.2 and 10.4 eV. The effect of a varying screening resulting from a change of electrolyte concentration was also accounted for.

(ii) In the second limiting case, one assumes that the transition dipole is represented by a point dipole. Detailed calculations of the dispersion shift have been performed, and they yield essentially the same functional dependence on photon energy for the dispersion spectrum as the model of complete charge separation. This is easily seen by comparing the experessions for the change of free energy resulting from dispersion for electronic polarization of the outer-sphere region according to the two models. The solvent is taken to be transparent to simplify matters. This change of free energy is

$$\Delta P_{CCS}^{out} = Ae^2/2a \tag{1}$$

according to the complete charge separation model, where e is the electronic charge;  $a = r_c + 2r_w$ , the terms  $r_c$  and  $r_w$  being the crystallographic radii of the ion being photoionized and water, respectively. One has

$$A = \begin{bmatrix} 1 - \varepsilon_{op}^{-1} \end{bmatrix} - \begin{bmatrix} 1 - (\varepsilon_{op}^{0})^{-1} \end{bmatrix}$$
$$= (\varepsilon_{op}^{0})^{-1} - \varepsilon_{op}^{-1}$$
(2)

where  $\varepsilon_{op}^0$  is the limiting value of the optical dielectric constant of the solvent in the near infrared, and  $\varepsilon_{op}$  the optical dielectric constant at the prevailing photon energy.

Conversely,

$$\Delta P_{pd}^{out} = B_{\mu}^{2}/a^{3} \tag{3}$$

according to the point dipole model, where  $\mu$  is the transition dipole and

$$B = (\epsilon_{op} -1)/(2\epsilon_{op} + 1) - (\epsilon_{op}^{0} - 1)/(2\epsilon_{op}^{0} + 1).$$
 (4)

Equations (3) and (4) follow directly from the Kirkwood treatment [7] of the solvation free energy of a dipole in a cavity of radius a surrounded by a continuous medium (cf. eq. (3) in [8]).

One deduces from (1) to (4)

$$\Delta P_{ccs}^{out}/\Delta P_{pd}^{out} = (K/2)(ea/\mu)^2$$
 (5)

with

$$K = A/B$$

$$= \left[2 + (\varepsilon_{op}^{0})^{-1}\right]\left[2 + (\varepsilon_{op})^{-1}\right]/3$$
(6)

The variation of K with  $\varepsilon_{op}$  shows how the two models differ in accounting for the effect of dispersion in the outer-sphere region for a transparent solvent. The range of interest in the present case is  $\varepsilon_{op}^0 < \varepsilon_{op} < 2.5$ , and one has K = 2.189 and 2.050 for  $\varepsilon_{op} = \varepsilon_{op}^0 = 1.777$  and  $\varepsilon_{op} = 2.5$ , respectively. Thus, K decreases only by 6.3 percent in a monotonic fashion as  $\varepsilon_{op}$  increases from 1.777 to 2.5 whereas A increases from 0 to 0.163. The two models therefore have essentially the same dependence on dielectric constant although the free energies of polarization have different analytical expressions.

The foregoing conclusion was reached for the outer-sphere region, but it should also be valid for the inner-sphere shell. The treatment of this region involves the polarizability of the solvent in both models, and dispersion is accounted for by the variation of this quantity with photon energy. Only the method of calculating the field acting on the solvent multipoles in the inner-sphere shell varies with the model, and the effect of dispersion is solely contained in the solvent polarizability.

Although the two models predict nearly the same functional dependence of dispersion spectra on photon energies, they yield very different values of the

dispersion shift. Thus, the ratio of (5) is ca.  $540/\mu^2$ , where  $\mu$  is in debyes, and the following data were selected: K=2.05 ( $\varepsilon_{\rm op}\approx2.5$ ) and a = 4.78 Å ( $r_{\rm C}=2$  Å). Comparison of the theoretical total interaction energy calculated in [4] with experiment in the 7 to 10 eV range shows that the model of complete charge separation yields dispersion shifts which are too high by a factor of ca. three in the absence of ionic screening. Conversely, the point dipole model, in view of the above treatment, should yield dispersion shifts which are too low by ca. one order of magnitude. This is not surprising because the model of a point dipole grossly underestimates the change of electric field around the species being photoionized. Conversely, this change of field is overestimated in the model of complete charge separation because of the finite distance between the charges thus involved. The model of complete charge separation seems preferable because it involves less cumbersome algebra than the treatment of the dipole model would entail without the simplifying assumption of a point dipole.

### 4. Conclusion

The interaction between solvent molecules and transition dipoles results in a measurable <u>nonequilibrium</u> contribution (dispersion shift) to the energetics of photoionization in the case of significant dielectric dispersion of the solvent at the prevailing photon energy. Two nonequilibrium contributions to the photoionization free energy of liquids and solutions therefore must be considered: (i) the <u>electronic</u> contribution discussed here (dispersion shift) and (ii) the free energy of <u>nuclear</u> reorganization [5]. The functional dependence of the dispersion shift on photon energy is accounted for by theory in agreement with experiment.

Acknowledgement

This work was supported by the Office of Naval Research.

### References

- [1] J. D. Macomber, The dynamics of spectroscopic transitions (Wiley, New York, 1976), pp. 95-108.
- [2] E. Merzbacher, Quantum mechanics (Wiley, New York, 1961), pp. 459-460.
- [3] P. Delahay and A. Dziedzic, J. Chem. Phys. 80 (1984) 5381.
- [4] P. Delahay and A. Dziedzic, J. Chem. Phys. 84 (1986) 936.
- [5] P. Delahay, in: Electron spectroscopy: Theory, techniques and applications, vol. 5, eds. C. R. Brundle and A. D. Baker (Academic Press, London, 1984), pp. 123-196.
- [6] J. J. Sakurai, Advanced quantum mechanics (Addison-Wesley, Reading, Mass., 1967), p. 37.
- [7] J. G. Kirkwood, J. Chem. Phys. 2 (1934) 351.
- [8] K. G. Spears, T. H. Gray and D. Huang, J. Phys. Chem. 90 (1986) 779.

### Captions to Figures

- Fig. 1. Dispersion spectra of 1 M aqueous solution of sodium azide with hydrogen (A) and argon (B) sources having the spectral outputs of fig. 2. Threshold energy of  $N_3^-$ ,  $E_t = 7.4$  eV. Very low output of the argon source above 9.0 eV (dashed portion of B).
- Fig. 2. Spectral outputs of hydrogen (A) and argon (B) sources in the experiments of fig. 1. Relative intensities of ca. 0.4 at the normalized highest peaks, the hydrogen source being the stronger.
- Fig. 3. Dispersion spectra of 1 M sodium iodide solutions in water (A) and glycerol (B). Dispersion spectrum of pure glycerol (C). Threshold energies,  $E_t \approx 7.4 \text{ eV}$  for  $I^-$  in water and glycerol;  $E_t \approx 8.7 \text{ eV}$  for pure glycerol;  $E_t = 10.06 \text{ eV}$  for pure water. All spectra obtained with hydrogen lamp.




FIG. I




FIG. 2




FIG. 3

## 01/1113/86/2

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 1911	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003

Dr. J. Driscoll Lockheed Palo Alto Research Laboratory 3251 Hanover Street Palo Alto, California 94304

Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602

Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125

Dr. J. J. Auborn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135

Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063

Dr. Roger Belt Litton Industries Inc. Airtron Division Morris Plains, NJ 07950

Dr. Ulrich Stimming Department of Chemical Engineering Columbia University New York, NY 10027 Dr. Manfred Breiter Institut fur Technische Elektrochemie Technischen Universitat Wien 9 Getreidemarkt, 1160 Wien AUSTRIA

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Laboratory
Livermore, California 94550

Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706

Dr. Steven Greenbaum
Department of Physics and Astronomy
Hunter College
695 Park Ave.
New York, NY 10021

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains Naval Weapons Support Center Electrochemical Power Sources Division Crane, Indiana 47522

S. Ruby DOE (STOR) Room 5E036 Forrestal Bldg., CE-14 Washington, D.C. 20595

Dr. A. J. Bard Department of Chemistry University of Texas Austin, Texas 78712

Dr. Janet Osteryoung Department of Chemistry State University of New York Buffalo, New York 14214

Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910

Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840

Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709

Dr. Aaron Fletcher Naval Weapons Center Code 3852 China Lake, California 93555 Dr. Michael J. Weaver Department of Chemistry Purdue University West Lafayette, Indiana 47907

Dr. R. David Rauh EIC Laboratories, Inc. Norwood, Massachusetts 02062

Dr. Aaron Wold Department of Chemistry Brown University Providence, Rhode Island 02192

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH UNITED KINGDOM

Dr. R. A. Osteryoung Department of Chemistry State University of New York Buffalo, New York 14214

Or. John Wilkes
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. D. Rolison Naval Research Laboratory Code 6171 Washington, D.C. 20375-5000

Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201

Dr. Edward M. Eyring Department of Chemistry University of Utah Salt Lake City, UT 84112

Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California

Dr. Hector D. Abruna Department of Chemistry Cornell University Ithaca, New York 14853

Dr. A. B. P. Lever Chemistry Department York University Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak Naval Ocean Systems Center Code 633, Bayside San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Micha Tomkiewicz Department of Physics Brooklyn College Brooklyn, New York 11210

Dr. Lesser Blum Department of Physics University of Puerto Rico Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II IBM Corporation 5600 Cottle Road San Jose, California 95193

Dr. Nathan Lewis Department of Chemistry Stanford University Stanford, California 94305 Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201

Dr. Alan Bewick Department of Chemistry The University of Southampton Southampton, SO9 5NH UNITED KINGDOM

Dr. E. Anderson NAVSEA-56Z33 NC #4 541 Jefferson Davis Highway Arlington, VA

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Or. Elton Cairns
Emergy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. Richard Pollard Department of Chemical Engineering University of Houston Houston, Texas 77004

Dr. M. Philpott IBM Corporation 5600 Cottle Road San Jose, California 95193

Dr. Donald Sandstrom Boeing Aerospace Co. P.O. Box 3999 Seattle, Washington 98124

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

Dr. Joel Harris Department of Chemistry University of Utah Salt Lake City, Utah 84112

Dr. Robert Somoano Jet Propulsion Laboratory California Institute of Technology Pasadena, California 91103

Dr. Johann A. Joebstl USA Mobility Equipment R&D Command DRDME-EC Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus NASA Headquarters M.S. RTS-6 Washington, D.C. 20546

Dr. Albert R. Landgrebe U.S. Department of Energy M.S. 68025 Forrestal Building Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin Department of Chemistry Texas A&M University College Station, Texas 77843

Dr. H. Tachikawa Department of Chemistry Jackson State University Jackson, Mississippi 39217

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl U.S. Department of Energy MS G-226 Washington, D.C. 20545

Dr. Edward Fletcher Department of Mechanical Engineering University of Minnesota Minneapolis, Minnesota 55455

Dr. John Fontanella Department of Physics U.S. Naval Academy Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson Syntheco, Inc. Rte 6 - Industrial Pike Road Gastonia, North Carolina 28052

Dr. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells Eltron Research Inc. 4260 Westbrook Drive, Suite 111 Aurora, Illinois 60505

Dr. C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

DL/1113/86/2

## ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. John Owen
Department of Chemistry and
Applied Chemistry
University of Salford
Salford M5 4WT ENGLAND

Dr. Boone Owens
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Menahem Anderman W.R. Grace & Co. Columbia, MD 20144

Dr. J. O. Thomas University of Uppsala Institute of Chemistry Box 531 S-751 21 Uppsala, Sweden

Dr. O. Stafsudd
Department of Electrical Engineering
University of California
Los Angeles, California 90024

and the transferior and th